


Inpatient exercises – capsular release & subacromial decompression




Early in-hospital exercises after capsular release or subacromial decompression.

Kieran Hirpara © ⓘ 4.0

These are the gentle exercises to begin in hospital after a capsular release or a subacromial decompression. For these operations the priority is to keep the shoulder moving freely.

Your exercises




Wrist movement

Keep your hand moving by bending your wrist forwards, backwards and side to side.

10 times, 3 times per day

Kieran Hirpara © ⓘ 4.0




Open and close hand

Keep your hand and fingers moving by opening and closing them, or by squeezing a stress ball.

10 times, 3 times per day

Kieran Hirpara © ⓘ 4.0




Elbow bends

Bend and straighten your elbow.

10 times, 3 times per day

Kieran Hirpara © ⓘ 4.0




Pendulums

This is a passive exercise. Lean forward and let your arm relax down. Use your body to move the arm gently either clockwise or anti-clockwise, along with forwards, backwards and side to side.

About 30 seconds each way, 3 times per day

Kieran Hirpara © ⓘ 4.0




Assisted forward flexion

Sitting on a chair and leaning forward, cradle your operated arm with the other arm and gently move it upwards in front of you. Lower it back down with the help of your non-operated arm. You could also lie on your back in bed and help the arm upwards if you prefer.

10 times, 3 times per day

Kieran Hirpara © ⓘ 4.0




Assisted abduction

Sitting on a chair and leaning forward, cradle the arm again and help it out to the side (like rocking a baby).

10 times, 3 times per day

Kieran Hirpara © ⓘ 4.0




External rotation

Sitting on a chair, only move your arm from where it sits in the sling to pointing straight in front of you. Don't go further outwards.

10 times, 3 times per day

Kieran Hirpara © ⓘ ⓘ 4.0



Lower trapezius setting

Squeeze your shoulder blades downwards and together.

Hold 5 seconds, 5 times, 3 times per day

Kieran Hirpara © ⓘ ⓘ 4.0




Upper trapezius stretch

Use your non-operated arm to bring your ear towards your shoulder, away from the operated side.

Hold 10 seconds, 3 times, 3 times per day

Kieran Hirpara © ⓘ ⓘ 4.0



Levator scapulae stretch

Use your non-operated arm to bring your nose towards your armpit area, away from the operated side.

Hold 10 seconds, 3 times, 3 times per day

Kieran Hirpara © ⓘ ⓘ 4.0

Begin these gentle exercises in hospital and continue them at home. They should be comfortable – ease back if pain increases.

CQ HAND + UPPER LIMB

Dr Kieran Hirpara – Specialist Orthopaedic Surgeon
Suite 2, Level 1, Mater Private Hospital Rockhampton, 31 Ward Street, The Range, QLD 4700
Phone 07 4863 6556 · office@cqupperlimb.com.au · cqupperlimb.com.au

Wearing your sling

- The sling is only to be worn for comfort, and should be left off as much as possible. You don't need to sleep in the sling.
- Move and use your arm as much as possible.
- It is advisable to wear the sling when out of the house, to prevent people knocking into your arm.
- Aim for physiotherapy sessions at least twice a week for the first six weeks.
- Use ice for pain relief if needed.
- When wearing your sling, relax your shoulder and let the sling take the weight of your arm.
- Take your painkillers before you do your exercises, and before your physiotherapy appointments.
- Unless you have arranged your own physiotherapy, an appointment has been made for you and is detailed in your discharge pack.
- If you have any problems, contact the rooms or let your physiotherapist know.

When you go home

Once you are home, your recovery continues with the full rehabilitation protocol for your operation: [capsular release](#); [subacromial decompression](#).