


Thoracic Outlet Syndrome

In thoracic outlet syndrome the nerves and vessels are squeezed at the base of the neck – often worse with the arm overhead.

Kieran Hirpara © ⓘ 4.0



What you're feeling

Thoracic outlet syndrome usually shows up as an arm that just isn't quite right. You might notice an ache through the shoulder, the side of the neck and down the arm, along with numbness, pins-and-needles or tingling, often in the ring and little fingers. The arm can feel heavy, weak or clumsy, so you fumble things or your grip tires quickly. A very typical clue is that the symptoms get **worse when your arm is up or overhead** (hanging out washing, reaching to a high shelf, driving, or carrying a bag on your shoulder) and ease again when you rest the arm down. Some people get headaches at the back of the head as well.

Less commonly, the problem is with the blood vessels rather than the nerves. If a vein is squeezed, the whole arm can suddenly swell, feel heavy and take on a bluish tinge. If an artery is squeezed (which is rare), the hand may go pale, cold and painful. These vascular patterns matter, because they need to be looked at quickly (see the last section).

What's actually happening

To get from your neck to your arm, the nerves and blood vessels have to pass through a narrow gap at the base of the neck, between your collarbone, your first rib, and the scalene muscles of the neck. This gap is the "thoracic outlet". When it becomes too tight, those structures get compressed, and that pressure is what produces the symptoms.

There are three types, depending on what is being squeezed. **Neurogenic TOS** (pressure on the nerves, the brachial plexus) is by far the commonest, making up the great majority of cases; this is the aching, numbness, tingling and weakness pattern. **Venous TOS** involves the main vein and causes the swollen, heavy, bluish arm, sometimes with a clot. **Arterial TOS** is rare and involves the artery, causing the pale, cold hand.

A few things make the space tighter. Some people are born with an **extra rib in the neck (a cervical rib)** or a tight fibrous band. Posture also matters: rounded, drooping shoulders close the space down, as do bulky or tight

neck and chest muscles, which is why the condition is more common in people who do repetitive overhead work or sport. A previous neck or collarbone injury can also be the trigger.

What we can do about it

At Mater Private Hospital Rockhampton, Dr Kieran Hirpara leads our approach to this condition. Patients are generally referred to our clinic by their GP; if a physiotherapist has suggested you see us, you will still need a referral from your GP in order to be eligible for the Medicare rebate, and begin with a thorough assessment to confirm the diagnosis. For long-standing symptoms, we usually start with non-operative care and consider surgery only if that does not provide enough relief.

The good news is that for the common, neurogenic type, the first-line treatment is physiotherapy, not surgery. A targeted programme to open up the thoracic outlet does the heavy lifting: correcting posture, strengthening the muscles that hold the shoulder blade back and down, releasing tight neck and chest muscles, and learning to avoid the arm positions that pinch. This settles symptoms for most people, and it's worth giving it a genuine, several-month effort.

Surgery is reserved for two situations. The first is the vascular types (venous or arterial), where decompression is usually needed and is the main treatment. The second is neurogenic TOS that hasn't settled despite a proper trial of physiotherapy and is genuinely interfering with life. The operation decompresses the outlet, typically by removing the first rib (and any extra cervical rib) and releasing the scalene muscles, to give the nerves and vessels room. It can be done through the armpit, above the collarbone, or with keyhole assistance. For selected patients it works well.

What to expect

TOS can be a tricky diagnosis. There's no single test that proves it. The diagnosis is mostly clinical: your story plus examination tests that put the arm in provoking positions. Scans and nerve studies are often used as much to rule out commoner causes (like carpal tunnel at the wrist, cubital tunnel at the elbow, or a neck/disc problem) as to confirm TOS itself. So part of the process is patiently sorting out where your symptoms are really coming from, and the answer isn't always clear-cut.

For the **neurogenic** type, most people improve with posture and physiotherapy and never need an operation, though it can take months and needs you to keep the habits going. When the **vascular** types or stubborn neurogenic cases do come to surgery, decompression relieves symptoms for most carefully selected patients, with recovery over several weeks. Because TOS sits at the most proximal point of the nerve pathway (higher up than carpal tunnel at the wrist or cubital tunnel at the elbow), it's sometimes just one piece of the puzzle, and the plan is tailored to you.

When to see someone

- A sudden, swollen, heavy or bluish arm: this can mean a vein is blocked or clotted and needs **urgent** assessment, the same day.
- A pale, cold, painful hand, or fingers that change colour: a possible arterial problem and a reason to seek **urgent** care.
- **Numbness, tingling, weakness or clumsiness** in the arm or hand that is persisting, worsening, or interfering with work, sleep or daily tasks.
- **Wasting (loss of muscle bulk) in the hand**, or symptoms that keep coming back despite a good trial of physiotherapy: worth a specialist opinion.
- If you've been told you have a cervical rib and are getting arm symptoms, worth having it assessed.

In more depth

This section goes further than you need for your own treatment decisions. Thoracic outlet syndrome is worth the extra reading because it splits into forms that could hardly be more different – one objectively demonstrable and occasionally limb-threatening, the other diagnosed largely on clinical judgement – and almost all the controversy attaches to the second.

THREE CONDITIONS SHARING ONE NAME

The thoracic outlet is the narrow space between the collarbone and first rib through which the nerves to the arm and the main artery and vein all pass. Compression there produces three distinct syndromes.

Venous and **arterial** TOS are vascular problems. They are uncommon, they are objectively demonstrable on imaging, and they can be urgent – a clotted subclavian vein producing a swollen blue arm, or arterial compression producing a cold hand and, in the worst case, emboli. Nobody disputes these exist or that they need treating.

Neurogenic TOS accounts for the great majority. Here the nerves are compressed, producing aching in the shoulder and arm, numbness typically along the little-finger side, and symptoms worsened by holding the arms overhead. The difficulty is that in most cases there is no test that confirms it – nerve conduction studies are frequently normal, and imaging shows no lesion.

Contemporary reviews respond to this by distinguishing vascular and **objectively verifiable** forms of neurogenic TOS from the rest [1], which is a careful way of acknowledging that the remainder rest on clinical judgement.

WHY THAT DISTINCTION DRIVES EVERYTHING

Where the diagnosis can be objectively demonstrated, treatment decisions follow the usual logic. Where it cannot, two failure modes arise: people with genuine compression go years without a diagnosis because their

tests are normal, and people whose arm pain has another cause undergo a substantial operation that cannot help them.

Both errors are well recognised, which is why the standard of proof before surgery is set higher here than for most compressions, and why a prolonged trial of physiotherapy – addressing posture, the scalene muscles and shoulder girdle mechanics – precedes any operation for the neurogenic form.

WHAT IS REMOVED, AND THE EVIDENCE ON HOW

Surgery decompresses the outlet by removing the first rib, dividing the scalene muscles, or both.

Comparing approaches across **2,010** patients, **Disabilities of the Arm, Shoulder and Hand and Derkash scores were significantly better** following **rib-sparing scalenectomy**, and **higher complication rates were reported after first rib resection** [2].

Against that, longer-standing series argue for the combined operation. Across **532** patients, a combined transaxillary first rib resection with transcervical scalenectomy was described as the most complete decompression with a better rate of symptom improvement and lower recurrence [3] – and recurrence was reported at **5% to 10%** when the combined procedure was used as the primary operation [4].

The two positions are not fully reconcilable from the published data. The more recent comparative analysis favours doing less; the large single-surgeon experience favours doing more, on the grounds that incomplete decompression is what drives recurrence. What both agree on is that recurrent TOS is a considerably harder problem than the first presentation, so getting the initial decision right matters more than the choice of technique.

THE PRACTICAL IMPLICATION

Given a diagnosis that is often clinical, an operation whose optimal extent is contested, and a recurrence problem that is difficult to salvage, this is a condition where a second opinion before surgery is reasonable rather than adversarial – particularly for the neurogenic form without objective confirmation.

REFERENCES

1. Panther EJ, Reintgen CD, Cueto RJ, Hao KA, Chim H, King JJ. Thoracic outlet syndrome: a review. *J Shoulder Elbow Surg.* 2022;31(11):e545-e561.
2. Blondin M, Garner GL, Hones KM, Nichols DS, Cox EA, Chim H. Considerations for surgical treatment of neurogenic thoracic outlet syndrome: a meta-analysis of patient-reported outcomes. *J Hand Surg Am.* 2023;48(6):585-94.
3. Atasoy E. A hand surgeon's further experience with thoracic outlet compression syndrome. *J Hand Surg Am.* 2010;35(9):1528-38.
4. Atasoy E. Recurrent thoracic outlet syndrome. *Hand Clin.* 2004;20(1):99-105.